

Expression and Purification of Soluble form of Human Parathyroid Hormone (rhPTH1-34) by Trx Tag in *E. coli*

Sanaz Yari, Farida Behzadian*, Hamideh Rouhani Nejad, Mohammadreza Masoumian, Mahdi Karimi

Department of Biosciences and Biotechnology, Malek-Ashtar University of Technology, Tehran, Iran.

Received: 11 Jul 2017

Revised : 20 Aug 2017

Accepted: 10 Sep 2017

Corresponding Author:

Farida Behzadian

Department of Biosciences and
Biotechnology, Malek-Ashtar
University of Technology, Tehran,
Iran.

Phone: +98-2122974603

E-mail: fbehzadian@yahoo.com

Abstract

Background: Parathyroid Hormone (PTH) is secreted by parathyroid glands and controls the level of calcium in bones and kidney. PTH is a small polypeptide with 84 amino acids, but the first 34 amino acids of which are enough for hormone biological activity and can be used in the treatment of Osteoporosis. The expression efficiency of recombinant human parathyroid hormone rhPTH (1-34) or Teriparatide using a cleavable fusion protein strategy was compared in two strains of *E. coli*.

Materials and Methods: A cassette was designed and fully synthesized for prokaryotic expression of rhPTH using pET system. From 5' to 3', the cassette consisted of: Trx tag to increase the solubility of protein, His tag for purification and detection of protein, enterokinase site to cleave all fusion moieties, and an optimized gene code for Teriparatide corresponding to the amino acid sequence of hPTH. This cassette was cloned into pET32a vector. The vector was simultaneously transformed and expressed in two different *E. coli* strains. The ability of strains for expression of this recombinant pharmaceutical was compared. Early expression was confirmed by SDS-PAGE and Western Blotting. The soluble fusion protein was harvested and purified by immobilized affinity chromatography. Then the fusion moiety was released from Teriparatide by enterokinase digestion.

Results: The fusion form of rhPTH was efficiently expressed in both *E. coli* strains. However, the percentage of the target protein to the total protein content in Rosetta-gami was more than its amount in BL21 (60 % vs 25%). The fusion protein was highly purified with Ni-NTA column. Up to 18.5 mg/ml of pure fusion protein has been obtained from 1-liter Rosetta-gami strain of *E. coli*. The pure Teriparatide was released by enterokinase digestion.

Conclusion: The pure rhPTH (1-34) produced here, could be the subject for biological activity and quality control assessments, and following formulation processing, it could be applied as a peptide drug in the treatment of Osteoporosis.

Keywords: Enterokinase; Fusion protein; Parathyroid hormone; BL21; Rosetta-gami

Please cite this article as: Yari S, Behzadian F, Rouhani Nejad H, Masoumian M, Karimi M. Expression and Purification of Soluble form of Human Parathyroid Hormone (rhPTH1-34) by Trx Tag in *E. coli*. Res Mol Med. 2017; 5 (3): 26-31

Introduction

Teriparatide or rhPTH (1-34) is the recombinant bioactive portion of human parathyroid hormone secreted by parathyroid glands. Parathyroid hormone (PTH) has 115 amino acids in its original form; the first 25 N-terminal amino acids are separated as a signal, the next 6 amino acids are separated during

secretion from Golgi, and finally, the mature form is secreted from parathyroid glands with 84 amino acids. This hormone affects the metabolism of calcium and phosphorus and acts directly on the kidneys and bone tissues(1). Studies show that all these 84 amino acids are not necessary for the

in two strains of *E. coli*.

Expression of rhPTH 1-34 in different *E. coli* hosts

The rhPTH-pET plasmid was transformed into *E. coli* Rosetta-gami and BL21 via transformation. Single colonies from transformed cells were used for inoculation of 5 ml pre-culture medium containing ampicillin (100mg/ml). The cultures were incubated for 24hr at 37 °C with the shake of 180 rpm. 250 µl of the pre-cultures were used for inoculation of 5 ml of LB medium (ampicillin 100mg/ml) at 37 °C until reaching the OD 600 of 0.7. For protein expression, Isopropyl β-D-1-thiogalactopyranoside (IPTG) was added to the final concentration of 1 mM. Expression was continued at 37 °C for 4hr and cells were harvested within desired intervals. Quantification of protein expression released in medium was carried out by image analysis of the SDS-PAGE in 14.5 percent scans using CLIQS (12).

Western blotting

Using specific antibody-antigen binding, recombinant protein can be detected. The samples were transferred from the gel onto nitrocellulose paper. After transfer, the paper was incubated for an hour in BSA 3% solution for blocking at room temperature with shake. Afterward, the paper was washed three times with TBS-T solution and was reacted with Anti-His6 (Roche Cat# 04905318001) diluted 1:6000 in TBS-T for 1hr at room temperature. After repeated washing of paper, it was incubated with the appropriate secondary antibody, anti-mouse IgG Peroxidase conjugate (Sigma Cat# A8924) diluted 1:10000 in TBS-T at the same condition. Then, rinsing was repeated and, finally, by adding a solution of 3, 3'-

diaminobenzidine (DAB) (Sigma Cat# D5637) in the darkness after about 3 minutes, the bands related to the recombinant protein were observed in the expected size.

Purification of recombinant PTH

The frozen biomass of *E. coli* was resuspended in washing buffer (50 mM Na₂ HPO₄/NaH₂ PO₄ -pH = 7.4). After the centrifuge (9000rpm for 5 min), the resulting cells were resuspended in lysis buffer (20 mM Tris, 0.5 M NaCl, 10 mM Imidazole, PMSF 1mM-pH=8). Then, sonication was performed in 100W at 4 °C (12 cycles, 10 seconds off and 15 seconds on) and centrifuged at 11,000 rpm. The supernatant containing the fusion protein was collected and applied to Ni-NTA resin (Chelating Sepharose Fast Flow). The column was equilibrated with buffer A (20mM Tris, 0.5 M NaCl, 20 mM Imidazole-pH=8) followed by washing with buffer A. The fusion protein was eluted with Linear gradient of Imidazole concentration by Gradient maker with buffer B (20 mM Tris, 0.5 M NaCl, 500mM Imidazole pH=8). Fractions containing His6-Thioredoxin-hPTH (1-34) fusion protein were pooled and were evaluated by SDS-PAGE gel.

Proteolytic cleavage of fusion protein with enterokinase

In order to cleave hPTH from its tag, the desalted fusion protein by dialysis process was digested with enterokinase enzyme (EK MAX, Invitrogen Co, USA) for 16hr at room temperature. After 16 hours, the results of the digestion were evaluated by SDS-PAGE gel.




Figure 2. A) Expression of rhPTH 1-34 in Rosetta- gami: Lane 1; before induction. Lanes 2, 3, 4, 5 and 6 were shown 30 min, 1hr, 2hr, 3hr and 4hr after induction, respectively; Lane 7, protein marker. B) Expression of rhPTH1-34 in BL21: lane 1 before induction; Lanes 2, 3, 4, 5 and 6 were shown 30 min, 1hr, 2hr, 3hr and 4hr after induction, respectively; Lane 7, protein marker.

Results

Construct and expression of recombinant hPTH with Trx tag

The gene sequence encoded for PTH (1-34) was inserted in the C terminal end of thioredoxin gene sequence. Between them, enterokinase specific recognition site of (Asp-Asp-Asp-Asp-Lys) was inserted as a specific recognition site. The construct was inserted in pET32a (+). The expression vector was transformed into two *E. coli* strains, including BL21 and Rosetta-gami.






Figure 3. Western blotting for rhPTH protein+Trx tag. Lane 1, a positive control (interferon alpha fused with His-tag). Lane 2, rhPTH protein+ Trx tag expression in the *Rosetta-gami*. Lane 3, Mw marker.



A



B

Figure 4. A) SDS-PAGE for fraction of outlet Ni-NTA chromatography. Lane 1, before induction. Lane 2, second supernatant. Lane 3, first supernatant. Lane 4, out let. Lane 5, fraction 7. Lane 6, fraction 8. Lane 7, fraction 14. Lane 8, fraction 15. Lane 9, fraction 16. Lane 10, fraction 18. Lane 11, fraction 19. Lane 12, fraction 20. Lane 13, fraction 26. Lane 14, fraction 27. Lane 15, MW. Lane 16, pooled fraction (11-16). Lane 17, pooled fraction (17-22). **B)** Profile of the eluted PTH protein from the Imidazole gradient. The best concentration can be seen in 14-22.

Cleavage of the rhPTH from Trx tag

The Trx tag (18 kDa) was detached from rhPTH through digestion with enterokinase (Figure 5). Two

bands were determined including Trx tag (18 kDa) and rhPTH (4 kDa) in the SDS-PAGE.

Confirm the expression of rhPTH protein+ Trx tag by Western blotting

Western blotting was performed before purification for confirmation of rhPTH1-34+Trx tag recombinant protein (22 kDa). Figure 3 is related to expression of target protein in Rosetta-gami.

Purification of rhPTH + Trx tag

The rhPTH + Trx tag proteins were applied to Ni-NTA chromatography. The purified protein was obtained by 70% concentration in the supernatant (Figure 4A). Pattern of the eluted PTH protein from the Imidazol gradient (Figure 4B).

Cleavage of the rhPTH from Trx tag

The Trx tag (18 kDa) was detached from rhPTH through digestion with enterokinase (Figure 5). Two bands were determined including Trx tag (18 kDa) and rhPTH (4 kDa) in the SDS-PAGE.




Figure 5. SDS-PAGE for digested fusion protein. Lane 1, Protein Ladder; Lanes 2 and 3, two same samples from digested protein rhPTH (4kD) and Trx tag (18kD); and lane 4 is undigested PTH fusion protein (22kD).

Discussion

Osteoporosis is a metabolic disorder due to bone tissue loses its strength and becomes highly sensitive to be broken. According to the reports, one in every two women and one in every four men over age 50 years old persons are suffering from this disease. The presence of this disease in Iran considerably increased from 15% of the population in 2001 to 34% in 2011. Based on the reports, 20% of Iranian women over 50 years old are affected by remarkable degrees of bone density reduction. Due to the significance of this disease and its high cost of therapy, using new treatment methods is clearly important.

In 2002, a human recombinant drug, Teriparatide (commercially named FORTEO), was produced by Lilly pharmaceutical company and released to the customer market for Osteoporosis therapy (13). This hormone is also the only effective anabolic drug against Osteoporosis that also is currently consumed in different countries such as Iran. The need for this drug in Iran will be increased in the near future because of the growing elderly population and life expectancy.

In 1991, Forsberg et al surveyed on expression of this hormone in BL21 *E. coli* that was fused to the carboxylic end of 15KD human IgG binding protein. They reported that, the final yield of this hormone was 5mg in each 1 liter of culture media (14). Oshika et al in 1994 studied the direct expression of this polypeptide in *E. coli* which was 27mg per liter based on their report (15). In the recent study that was done in 2017 by Bakhtiari et al in Iran, expression of Teriparatide was approximately 43% of whole protein expression that was less than the present study (16).

Teriparatide is a small polypeptide that has been mostly studied in fusion with other proteins like NUS

A, IgG and MBP for prevention of digestion by host proteases. In our study, Trx fused expression of Teriparatide was investigated in two hosts, including Rosetta-gami and BL21 (17). In addition, this molecular tag increased the expression level of Teriparatide by causing soluble expression of this product instead of the inclusion form, as well as, preventing deformation of protein structure in the cytoplasm by effecting on disulfide bindings leading to reduction in downstream procedures and hence save a lot in costs. Our results showed that the Teriparatide expression was 45% of the total protein in Rosetta-gami that was more in comparison with the BL21 and other surveys.

Using the molecular labels was almost unavoidable in the expression process of many recombinant proteins, while, the presence of some excess factors in the final products was used of a drug is impossible. Thus, detachment of adjunct factors of the main product after purification is important in recombinant drug producing process. Enterokinase enzymes, by cutting the specific sites with no addition of any amino acids to the product and without harm or effect to humans, are one of the most common ways for detachment of the protein labels. Therefore, enterokinase was used to obtain 4 kDa rhPTH purified product(18).

According to presented method in this study, the efficiency of the fused form of rhPTH expression is equal to 26 mg/liter of culture media of which 70% was purified. This yield is a suitable output compared to the other investigations and can be improved by using fermenter system and can be used as a drug candidate for Osteoporosis therapy following the functional evaluation of purified protein in both in vitro and in vivo. Since this product was obtained in soluble form and in proper concentration by using the mentioned procedure, it can be applied as an appropriate way for producing this polypeptide.

Acknowledgments

This article was supported by research center for Biosciences and Biotechnology, Malek Ashtar University of Tehran.

Author contributions

FB and HR designed and directed the project, SY and MK performed the experiments, SY and FB wrote the article.

Conflict of Interest

There is no Conflict of Interest.

Support/Funding

This work was supported by Malek-Ashtar University.

References

1. Jin L BS, Chandrasekhar S. crystal structure of human parathyroid hormone 1-34 at 0.9-A resolution. *J Biol Chem.* 2000; 275:27238-44. PMID: 10837469
2. Schluter KD WM, Piper HM. parathyroid hormone induces protein kinase C but not adenylate cyclase in adult cardiomyocytes and regulates cyclic AMP levels via protein kinase C dependent. *Biochem J.* 1995; 310:439-44. PMID: 7654180
3. Sugiyama T, Torio T, Miyajima T, Kim YT, Oda H. Romosozumab and blosozumab: alternative drugs of mechanical strain-related stimulus toward a cure for osteoporosis. *Frontiers Endocrinol.* 2015; 6:54.
4. Ebina K, Hashimoto J, Kashii M, Hirao M, Kaneshiro S, Noguchi T, et al. The effects of switching daily teriparatide to oral bisphosphonates or denosumab in patients with primary osteoporosis. *J Bone Miner Metab.* 2017; 35(1):91-8. PMID: 26762133
5. Partridge NC, Li X, Qin L. Understanding Parathyroid Hormone Action. *Ann N Y Acad Sci.* 2006; 1068: 187-93. PMID: 16831918
6. Vad R NE, Dahl LA. Engineering of a *Pichia pastoris* expression system for secretion of high amounts of intact human parathyroid hormone. *J Biotechnol.* 2005; 116(3):251-60. PMID: 15707686
7. Murby M CL, Nilsson J. Stabilization of recombinant proteins from proteolytic degradation in *E. coli*. *Biotechnol Appl Biochem.* 1991; 14(3): 336-46. PMID: 1777118
8. Suzuki Y, Yabuta M, Ohsuye K. High level production of recombinant human parathyroid hormone 1-34. *Appl Environ Microb.* 1998; 64(2):526-9. PMID: 9464388
9. Fu XY, Tong WY, Wei DZ. Extracellular production of human parathyroid hormone as a thioredoxin fusion form in *E. coli* by chemical permeabilization combined with heat treat. *Biotech Prog* 2005; 21(5):1429-35. PMID: 16209546
10. He X, Li L, Xu H, Xi J, Cao X, Rong S, et al. A rice jacalin-related mannose-binding lectin gene, OsJRL, enhances *Escherichia coli* viability under high salinity stress and improves salinity tolerance of rice. *Plant Biol.* 2017; 19(2):257-67.
11. Arnau J, Lauritzen C, Petersen GE, Pedersen J. Current strategies for the use of affinity tags and tag removal for the purification of recombinant proteins. *Protein Expr Purif.* 2006; 48(1): 1-13. PMID: 16427311
12. Heras J, Dominguez C, Mata E, Pascual V, Lozano C, Torres, and Zarazaga M. GelJ – a tool for analyzing DNA fingerprint gel images. *BMC Bioinformatics.* 2015; 16: 270. PMID: 26307353
13. Grover HS, Luthra S, Maroo S. Teriparatide: A Novel Means to Ultimately Achieve True Regeneration. *J Clin Diagn Res.* 2013; 7(8): 1820-3. PMID: 24086927
14. Forsberg G, Brobjer M, Holmgren E, Bergdahl K, Persson P, Gautvik KM, et al. Thrombin and H64A subtilisin cleavage of fusion proteins for preparation of human recombinant parathyroid hormone. *J Protein Chem.* 1991; 10(5): 517-26. PMID: 1799410
15. Liu Q, Lin J, Liu M, Tao X, Wei D, Ma X, et al. Large scale preparation of recombinant human parathyroid hormone 1–84 from *Escherichia coli*. *Protein Expr Purif.* 2007; 54(2): 212-9. PMID: 17449274
16. Bakhtiari N Amini Bayat Z, Sagharidouz S, Vaez M. Overexpression of Recombinant Human Teriparatide, rhPTH (1-34) in *Escherichia coli*: an innovative gene fusion approach. *Avicenna J Med Biotechnol.* 2017; 9(1): 19-22. PMID: 28090276
17. Hamedifar H, Salamat F, Saffarion M, Ghiasi M, Hosseini A, Lahiji H, et al. A novel approach for high level expression of soluble recombinant human parathyroid hormone (rhPTH 1-34) in *E. coli*. *Avicenna J Med Biotechnol.* 2013; 5(3): 193-201. PMID: 23919123
18. Edward RL, Elizabeth AD, Charlotte K, Kathleen LG, Paul FS, John MM. A thioredoxin gene fusion expression system that circumvents inclusion body formation in the *E.coli* cytoplasm. *Biotechnol.* 1993; 11: 187-93.